

# EMERGENCE OF SPACETIME

B.G. Sidharth

International Institute for Applicable Mathematics & Information Sciences  
Hyderabad (India) & Udine (Italy)

B.M. Birla Science Centre, Adarsh Nagar, Hyderabad - 500 063 (India)

## Abstract

Starting from a background Zero Point Field (or Dark Energy) we show how an array of oscillators at the Planck scale leads to the formation of elementary particles and spacetime and also to a cosmology consistent with latest observations.

## 1 Introduction

Our starting point is the scenario in which there is particulate condensation in the all pervading Zero Point Field. This would be at the Compton length, which includes the Planck length as a special case. Indeed this was Einstein's belief. As Wilzeck put it, "Einstein was not satisfied with the dualism. He wanted to regard the fields, or ethers, as primary. In his later work, he tried to find a unified field theory, in which electrons (and of course protons, and all other particles) would emerge as solutions in which energy was especially concentrated, perhaps as singularities. But his efforts in this direction did not lead to any tangible success."

Let us see how this can happen. In the words of Wheeler, 'From the zero-point fluctuations of a single oscillator to the fluctuations of the electromagnetic field to geometrodynamical fluctuations is a natural order of progression...'

Let us consider, following Wheeler a harmonic oscillator in its ground state. The probability amplitude is

$$\psi(x) = \left(\frac{m\omega}{\pi\hbar}\right)^{1/4} e^{-(m\omega/2\hbar)x^2}$$

for displacement by the distance  $x$  from its position of classical equilibrium. So the oscillator fluctuates over an interval

$$\Delta x \sim (\hbar/m\omega)^{1/2}$$

The electromagnetic field is an infinite collection of independent oscillators, with amplitudes  $X_1, X_2$  etc. The probability for the various oscillators to have amplitudes  $X_1, X_2$  and so on is the product of individual oscillator amplitudes:

$$\psi(X_1, X_2, \dots) = \exp[-(X_1^2 + X_2^2 + \dots)]$$

wherein there would be a suitable normalization factor. This expression gives the probability amplitude  $\psi$  for a configuration  $B(x, y, z)$  of the magnetic field that is described by the Fourier coefficients  $X_1, X_2, \dots$  or directly in terms of the magnetic field configuration itself by

$$\psi(B(x, y, z)) = P \exp \left( - \int \int \frac{\mathbf{B}(\mathbf{x}_1) \cdot \mathbf{B}(\mathbf{x}_2)}{16\pi^3 \hbar c r_{12}^2} d^3 x_1 d^3 x_2 \right).$$

$P$  being a normalization factor. Let us consider a configuration where the magnetic field is everywhere zero except in a region of dimension  $l$ , where it is of the order of  $\sim \Delta B$ . The probability amplitude for this configuration would be proportional to

$$\exp[-(\Delta B)^2 l^4 / \hbar c]$$

So the energy of fluctuation in a region of length  $l$  is given by finally

$$B^2 \sim \frac{\hbar c}{l^4}$$

In the above if  $l$  is taken to be the Compton wavelength of a typical elementary particle, then we recover its energy  $mc^2$ , as can be easily verified. In Quantum Gravity as well as in Quantum Super String Theory, we encounter phenomena at a minimum scale. It is well known, and this was realized by Planck himself, that there is an absolute minimum scale in the universe, and this is,

$$\begin{aligned} l_P &= \left( \frac{\hbar G}{c^3} \right)^{\frac{1}{2}} \sim 10^{-33} \text{ cm} \\ t_P &= \left( \frac{\hbar G}{c^5} \right)^{\frac{1}{2}} \sim 10^{-42} \text{ sec} \end{aligned} \tag{1}$$

Yet what we encounter in the real world is, not the Planck scale, but the elementary particle Compton scale. The explanation given for this is that the very high energy Planck scale is moderated by the Uncertainty Principle. The question which arises is, exactly how does this happen? We will now present an argument to show how the Planck scale leads to the real world Compton scale, via fluctuations and the modification of the Uncertainty Principle. We note that (1) defines the absolute minimum physical scale [1, 2, 3, 4]. Associated with (1) is the Planck mass

$$m_P \sim 10^{-5} gm \quad (2)$$

There are certain interesting properties associated with (1) and (2).  $l_P$  is the Schwarzschild radius of a black hole of mass  $m_P$  while  $t_P$  is the evaporation time for such a black hole via the Beckenstein radiation [5]. Interestingly  $t_P$  is also the Compton time for the Planck mass, a circumstance that is symptomatic of the fact that at this scale, electromagnetism and gravitation become of the same order [6]. Indeed all this fits in very well with Rosen's analysis that such a Planck scale particle would be a mini universe [7, 8]. We will now invoke a time varying gravitational constant discussed extensively in ref.[6].

$$G \approx \frac{lc^2}{m\sqrt{N}} \propto (\sqrt{N}t)^{-1} \propto T^{-1} \quad (3)$$

which resembles the Dirac cosmology and features in the author's successful 1997 model, in which (3) arises due to the fluctuation in the particle number [9, 10, 11, 12, 6]. In (3)  $m$  and  $l$  are the mass and Compton wavelength of a typical elementary particle like the pion while  $N \sim 10^{80}$  is the number of elementary particles in the universe, and  $T$  is the age of the universe.

In this scheme wherein (3) follows from the theory, we use the fact that given  $N$  particles, the fluctuation in the particle number is of the order  $\sqrt{N}$ , while a typical time interval for the fluctuations is  $\sim \hbar/mc^2$ , the Compton time. We will come back to this point later. So anticipating later work we have

$$\frac{dN}{dt} = \frac{\sqrt{N}}{\tau}$$

whence on integration we get,

$$T = \frac{\hbar}{mc^2} \sqrt{N}$$

and we can also deduce its spatial counterpart,  $R = \sqrt{N}l$ , which is the well known empirical Eddington formula.

Equation (3) which is an order of magnitude relation is consistent with observation [13, 14] while it may be remarked that the Dirac cosmology itself has inconsistencies.

Substitution of (3) in (1) yields

$$\begin{aligned} l &= N^{\frac{1}{4}}l_P, \\ t &= N^{\frac{1}{4}}t_P \end{aligned} \tag{4}$$

where  $t$  as noted is the typical Compton time of an elementary particle. We can easily verify that (4) is correct. It must be stressed that (4) is not a fortuitous empirical coincidence, but rather is a result of using (3), which again as noted, can be deduced from theory.

(4) can be rewritten as

$$\begin{aligned} l &= \sqrt{n}l_P \\ t &= \sqrt{n}t_P \end{aligned} \tag{5}$$

wherein we have used (1) and (3) and  $n = \sqrt{N}$ .

We will now compare (5) with the well known relations, referred to earlier,

$$R = \sqrt{N}l \quad T = \sqrt{N}t \tag{6}$$

The first relation of (6) is the well known Weyl-Eddington formula referred to while the second relation of (6) is given also on the right side of (3). We now observe that (6) can be seen to be the result of a Brownian Walk process,  $l, t$  being typical intervals between "steps" (Cf.[6, 15, 16]). We demonstrate this below after equation (8). On the other hand, the typical intervals  $l, t$  can be seen to result from a diffusion process themselves. Let us consider the well known diffusion relation,

$$(\Delta x)^2 \equiv l^2 = \frac{\hbar}{m}t \equiv \frac{\hbar}{m}\Delta t \tag{7}$$

(Cf.[15],[17]-[20]). What is being done here is that we are modeling fuzzy spacetime by a double Wiener process to be touched upon later, which leads to (7). This will be seen in more detail, below.

Indeed as  $l$  is the Compton wavelength, (7) can be rewritten as the Quantum Mechanical Uncertainty Principle

$$l \cdot p \sim \hbar$$

at the Compton scale (Cf. also [21]) (or even at the de Broglie scale). What (7) shows is that a Brownian process defines the Compton scale while (6) shows that a Random Walk process with the Compton scale as the interval defines the length and time scales of the universe itself (Cf.[16]). Returning now to (5), on using (2), we observe that in complete analogy with (7) we have the relation

$$(\Delta x)^2 \equiv l_P^2 = \frac{\hbar}{m_P} t_P \equiv \frac{\hbar}{m_P} \Delta t \quad (8)$$

We can now argue that the Brownian process (8) defines the Planck length while a Brownian Random Walk process with the Planck scale as the interval leads to (5), that is the Compton scale.

To see all this in greater detail, it may be observed that equation (8) (without subscripts)

$$(\Delta x)^2 = \frac{\hbar}{m} \Delta t \quad (9)$$

is the same as the equation (7), indicative of a double Wiener process. Indeed as noted by several scholars, this defines the fractal Quantum path of dimension 2 (rather than dimension 1) (Cf.e.g. ref.[18]).

Firstly it must be pointed out that equation (9) defines a minimum space time unit - the Compton scale  $(l, t)$ . This follows from (9) if we substitute into it  $\langle \frac{\Delta x}{\Delta t} \rangle_{max} = c$ . If the mass of the particle is the Planck mass, then this Compton scale becomes the Planck scale.

Let us now consider the distance traversed by a particle with the speed of light through the time interval  $T$ . The distance  $R$  covered would be

$$\int dx = R = c \int dt = cT \quad (10)$$

by conventional reasoning. In view of the equation (8), however we would have to consider firstly, the minimum time interval  $t$  (Compton or Planck time), so that we have

$$\int dt \rightarrow nt \quad (11)$$

Secondly, because the square of the space interval  $\Delta x$  (rather than the interval  $\Delta x$  itself as in conventional theory) appears in (8), the left side of (10) becomes, on using (11)

$$\int dx^2 \rightarrow \int (\sqrt{n} dx)(\sqrt{n} dy) \quad (12)$$

Whence for the linear dimension  $R$  we would have

$$\sqrt{n}R = nct \quad \text{or} \quad R = \sqrt{n}l \quad (13)$$

Equation (12) brings out precisely the fractal dimension  $D = 2$  of the Brownian path while (13) is identical to (4) or (6) (depending on whether we are dealing with minimum intervals of the Planck scale or Compton scale of elementary particles). Apart from showing the Brownian character linking equations (4) and (8), incidentally, this also provides the justification for what has so far been considered to be a mysterious large number coincidence viz. the Eddington formula (6).

There is another way of looking at this. It is well known that in approaches like that of the author or Quantum Super String Theory, at the Planck scale we have a non commutative geometry encountered earlier [22, 23] Indeed this follows without recourse to Quantum Super Strings, merely by the fact that  $l_P, t_P$  are the absolute minimum space time intervals as shown by Snyder [8]. as we saw earlier.

The non commutative geometry as is known, is symptomatic of a modified uncertainty principle at this scale [24]-[29]

$$\Delta x \approx \frac{\hbar}{\Delta p} + l_P^2 \frac{\Delta p}{\hbar} \quad (14)$$

The relation (14) would be true even in Quantum Gravity. The extra or second term on the right side of (14) expresses the well known duality effect - as we attempt to go down to the Planck scale, infact we are lead to the larger scale. The question is, what is this larger scale? If we now use the fact that  $\sqrt{n}$  is the fluctuation in the number of Planck particles (exactly as  $\sqrt{N}$  was the fluctuation in the particle number as in (3)) so that  $\sqrt{n}m_{PC} = \Delta p$  is the fluctuation or uncertainty in the momentum for the second term on the right side of (14), we obtain for the uncertainty in length,

$$\Delta x = l_P^2 \frac{\sqrt{n}m_{PC}}{\hbar} = l_P \sqrt{n}, \quad (15)$$

We can easily see that (15) is the same as the first relation of (5). The second relation of (5) follows from an application of the time analogue of (14).

Thus the impossibility of going down to the Planck scale because of (14), manifests itself in the fact that as we attempt to go down to the Planck scale, we infact end up at the Compton scale. This is how the Compton

scale is encountered in real life.

Interestingly while at the Planck length, we have a life time of the order of the Planck time, as noted above it is possible to argue on the other hand that with the mass and length of a typical elementary particle like the pion, at the Compton scale, we have a life time which is the age of the universe itself as shown by Sivaram [5, 30].

Interestingly also Ng and Van Dam deduce the relations like [31]

$$\delta L \leq (Ll_P^2)^{1/3}, \delta T \leq (Tt_P^2)^{1/3} \quad (16)$$

where the left side of (16) represents the uncertainty in the measurement of length and time for an interval  $L, T$ . We would like to point out that if in the above we use for  $L, T$ , the size and age of the universe, then  $\Delta L$  and  $\Delta T$  reduce to the Compton scale  $l, t$ .

In conclusion, Brownian double Wiener processes and the modification of the Uncertainty Principle at the Planck scale lead to the physical Compton scale.

## 2 The Universe as Planck Oscillators

In the previous section, we had argued that a typical elementary particle like a pion could be considered to be the result of  $n \sim 10^{40}$  evanescent Planck scale particles. We will return to this line of thinking again. The argument was based on random motions and also on the modification to the Uncertainty Principle. We will now consider the problem from a different point of view, which not only reconfirms the above result, but also enables an elegant extension to the case of the entire universe itself. Let us consider an array of  $N$  particles, spaced a distance  $\Delta x$  apart, which behave like oscillators, that is as if they were connected by springs. We then have [32, 33]

$$r = \sqrt{N\Delta x^2} \quad (17)$$

$$ka^2 \equiv k\Delta x^2 = \frac{1}{2}k_B T \quad (18)$$

where  $k_B$  is the Boltzmann constant,  $T$  the temperature,  $r$  the extent and  $k$  is the spring constant given by

$$\omega_0^2 = \frac{k}{m} \quad (19)$$

$$\omega = \left( \frac{k}{m} a^2 \right)^{\frac{1}{2}} \frac{1}{r} = \omega_0 \frac{a}{r} \quad (20)$$

We now identify the particles with Planck masses, set  $\Delta x \equiv a = l_P$ , the Planck length. It may be immediately observed that use of (19) and (18) gives  $k_B T \sim m_P c^2$ , which ofcourse agrees with the temperature of a black hole of Planck mass. Indeed, as noted, Rosen had shown that a Planck mass particle at the Planck scale can be considered to be a universe in itself. We also use the fact alluded to that a typical elementary particle like the pion can be considered to be the result of  $n \sim 10^{40}$  Planck masses. Using this in (17), we get  $r \sim l$ , the pion Compton wavelength as required. Further, in this latter case, using (48) and the fact that  $N = n \sim 10^{40}$ , and (18), i.e.  $k_B T = kl^2/N$  and (19) and (20), we get for a pion, remembering that  $m_P^2/n = m^2$ ,

$$k_B T = \frac{m^3 c^4 l^2}{\hbar^2} = mc^2,$$

which of course is the well known formula for the Hagedorn temperature for elementary particles like pions. In other words, this confirms the conclusions in the previous section, that we can treat an elementary particle as a series of some  $10^{40}$  Planck mass oscillators. However it must be observed from (18) and (19), that while the Planck mass gives the highest energy state, an elementary particle like the pion is in the lowest energy state. This explains why we encounter elementary particles, rather than Planck mass particles in nature. Infact as already noted [6], a Planck mass particle decays via the Bekenstein radiation within a Planck time  $\sim 10^{-42} \text{secs}$ . On the other hand, the lifetime of an elementary particle would be very much higher.

In any case the efficacy of our above oscillator model can be seen by the fact that we recover correctly the masses and Compton scales in the order of magnitude sense and also get the correct Bekenstein and Hagedorn formulas as seen above, and get the correct estimate of the mass of the universe itself, as will be seen below.

Using the fact that the universe consists of  $N \sim 10^{80}$  elementary particles like the pions, the question is, can we think of the universe as a collection of  $nN$  or  $10^{120}$  Planck mass oscillators? This is what we will now show. Infact if we use equation (17) with

$$\bar{N} \sim 10^{120},$$



we can see that the extent  $r \sim 10^{28} \text{cms}$  which is of the order of the diameter of the universe itself. Next using (20) we get

$$\hbar\omega_0^{(min)} \langle \frac{l_P}{10^{28}} \rangle^{-1} \approx m_P c^2 \times 10^{60} \approx M c^2 \quad (21)$$

which gives the correct mass  $M$ , of the universe which in contrast to the earlier pion case, is the highest energy state while the Planck oscillators individually are this time the lowest in this description. In other words the universe itself can be considered to be described in terms of normal modes of Planck scale oscillators.

We will return to these considerations later: this and the preceeding considerations merely set the stage.

In the above cosmology of fluctuations, our starting point was the creation of  $\sqrt{N}$  particles within the minimum time interval, a typical elementary particle Compton time  $\tau$ . A rationale for this, very much in the spirit of the condensation of particles from a background Zero Point Field as discussed can also be obtained in terms of a phase transition from the Zero Point Field or Quantum Vacuum as we will see in the sequel. In this case, particles are like the Benard cells which form in fluids, as a result of a phase transition. While some of the particles or cells may revert to the Zero Point Field, on the whole there is a creation of  $\sqrt{N}$  of these particles. If the average time for the creation of the  $\sqrt{N}$  particles or cells is  $\tau$ , then at any point of time where there are  $N$  such particles, the time elapsed, in our case the age of the universe, would be given by (6) (Cf. [34, 16]). While this is not exactly the Big Bang scenario, there is nevertheless a rapid creation of matter from the background Quantum Vacuum or Zero Point Field. Thus over  $10^{40}$  particles would have been created within a fraction of a second.

In any case when  $\tau \rightarrow 0$ , we recover the Big Bang scenario with a singular creation of matter, while when  $\tau \rightarrow$  Planck time we recover the Prigogine Cosmology (Cf.[8] for details). However in neither of these two limits we can deduce all the above consistent with observation large number relations which therefore have to be branded as accidents.

The above cosmological model is related to the fact that there are minimum space time intervals  $l, \tau$ . Indeed in this case as we saw there is an underlying non commutative geometry of spacetime [35, 23, 36] given by

$$[x, y] \approx 0(l^2), [x, p_x] = i\hbar[1 + \beta l^2], [t, E] = i\hbar[1 + \gamma \tau^2] \quad (22)$$

Interestingly (22) implies as we saw, modification to the usual Uncertainty Principle. (This in turn has also been interpreted in terms of a variable speed of light cosmology [37, 38, 39]).

The relations (22), lead to the modified Uncertainty relation

$$\Delta x \sim \frac{\hbar}{\Delta p} + \alpha' \frac{\Delta p}{\hbar} \quad (23)$$

(23) appears also in Quantum Super String Theory and is related to the well known Duality relation

$$R \rightarrow \alpha'/R$$

(Cf.[26, 3]). In any case (23) is symptomatic of the fact that we cannot go down to arbitrarily small space time intervals. We now observe that the first term of (23) gives the usual Uncertainty relation. In the second term, we write  $\Delta p = \Delta N mc$ , where  $\Delta N$  is the Uncertainty in the number of particles, in the universe. Also  $\Delta x = R$ , the radius of the universe where

$$R \sim \sqrt{N}l,$$

the famous Eddington relationship (9). It should be stressed that the otherwise empirical Eddington formula, arises quite naturally in a Brownian characterisation of the universe as has been pointed out in the previous Chapter (Cf. for example ref.[32]). Put simply (9) is the Random Walk equation.

We now get back,

$$\Delta N = \sqrt{N}$$

This is the uncertainty in the particle number, we used earlier. Substituting this in the time analogue of the second term of (23), we immediately get,  $T$  being the age of the universe,

$$T = \sqrt{N}\tau$$

which is equation (6). So, our cosmology is self consistent with the modified relation (23). The fluctuational effects are really couched in the modification of the Heisenberg Principle, as given in (23).

Interestingly these minimum space time considerations can be related to the Feynmann-Wheeler Instantaneous Action At a Distance formulation (Cf.[40,

41, 42]), a point which we shall elaborate further in the sequel.

We finally remark that relations like (22) and (23), which can also be expressed in the form,  $a$  being the minimum length,

$$[x, p_x] = i\hbar[1 + \left(\frac{a}{\hbar}\right)^2 p^2]$$

(and can be considered to be truncated from a full series on the right hand side (Cf. [22]), could be deduced from the rather simple model of a lattice - a one dimensional lattice for simplicity. In this case we will have (Cf.[8])

$$[x, p_x] = i\hbar \cos\left(\frac{p}{\hbar}a\right),$$

where  $a$  is the lattice length,  $l$  the Compton length in our case. The energy time relation now leads to a correction to the mass energy formula, viz

$$E = mc^2 \cos(kl), k \equiv p/\hbar$$

This is the contribution of the extra term in the Uncertainty Principle.

As noted the Planck scale is an absolute minimum scale in the universe. We argued that with the passage of time the Planck scale would evolve to the present day elementary particle Compton scale. To recapitulate: We have by definition

$$\hbar G/c^3 = l_P^2$$

where  $l_P$  is the Planck length  $\sim 10^{-33} \text{ cms}$ . If we use  $G$  from (3) in the above we will get

$$l = N^{1/4} l_P \tag{24}$$

Similarly we have

$$\tau = N^{1/4} \tau_P \tag{25}$$

In (24) and (25)  $l$  and  $\tau$  denote the typical elementary particle Compton length and time scale, and  $N$  is the number of such elementary particles in the universe.

We could explain these equations in terms of the Benard cell like elementary particles referred to above. This time there are a total of  $n = \sqrt{N}$  Planck particles and (24) and (25) are the analogues of equations (6) in the context of the formation of such particles. Indeed as we saw a Planck mass,

$m_P \sim 10^{-5} gms$ , has a Compton life time and also a Bekenstein Radiation life time of the order of the Planck time. These spacetime scales are much too small and we encounter much too large energies from the point of view of our experimental constraints. As noted in the previous chapter our observed scale is the Compton scale, in which Planck scale phenomena are moderated. In any case it can be seen from the above that as the number of particles  $N$  increases, the scale evolves from the Planck to the Compton scale.

So, the scenario which emerges is, that as the universe evolves, Planck particles form the underpinning for elementary particles, which in turn form the underpinning for the universe by being formed continuously.

This can be confirmed by the following argument: We can rewrite (24) as

$$l = \nu' \sqrt{T} \quad (26)$$

$$\nu' = l_P / \sqrt{\tau} \approx \hbar / m_P$$

Equation (26) as we saw earlier, is identical to the Brownian diffusion process which is infact the underpinning for equations like (??) or (??), except that this time we have the same Brownian Theory operating from the Planck scale to the Compton scale, instead of from the Compton scale to the edge of the universe as seen above (Cf. also [32, ?]).

Interestingly, let us apply the above scenario of  $\sqrt{n}$  Planck particles forming an elementary particle, to the extra term of the modified Uncertainty Principle (23), as we did earlier in this section in (iv). Remembering that  $\alpha' = l_P^2$  in the theory, and  $\Delta p = N^{1/4} m_P c$ , in this case, we get, as  $\Delta x = l$ ,

$$l = N^{1/4} l_P,$$

which will be recognized as (24) itself! Thus once again we see how the above cosmology is consistently tied up with the non commutative space time expressed by equations (22) or (23).

It may be mentioned that, as indeed can be seen from (24) and (25), in this model, the velocity of light remains constant.

We would now like to comment further upon the Compton scale and the fluctuational creation of particles alluded to above. In this case particles

are being produced out of a background Quantum Vacuum or Zero Point Field which is pre space time. First a Brownian process alluded to above defines the Planck length while a Brownian random process with the Planck scale as the fundamental interval leads to the Compton scale (Cf. also ref.[4]).

This process is a phase transition, a critical phenomenon. To see this briefly, let us start with the Landau-Ginsburg equation [27]

$$-\frac{\hbar^2}{2m}\nabla^2\psi + \beta|\psi|^2\psi = -\alpha\psi \quad (27)$$

Here  $\hbar$  and  $m$  have the same meaning as in usual Quantum Theory. It is remarkable that the above equation (27) is identical with a similar Schrodinger like equation based on amplitudes which is discussed in [8], where moreover  $|\psi|^2$  is proportional to the mass (or density) of the particle (Cf. ref.[8] for details). The equation in question is,

$$i\hbar\frac{\partial\psi}{\partial t} = \frac{-\hbar^2}{2m'}\frac{\partial^2\psi}{\partial x^2} + \int \psi^*(x')\psi(x)\psi(x')U(x')dx', \quad (28)$$

In (28),  $\psi(x)$  is the probability of a particle being at the point  $x$  and the integral is over a region of the order of the Compton wavelength. From this point of view, the similarity of (28) with (27) need not be surprising considering also that near critical points, due to universality diverse phenomena like magnetism or fluids share similar mathematical equations. Equation (28) was shown to lead to the Schrodinger equation with the particle acquiring a mass (Cf.also ref.[28]).

Infact in the Landau-Ginsburg case the coherence length is given by

$$\xi = \left(\frac{\gamma}{\alpha}\right)^{\frac{1}{2}} = \frac{h\nu_F}{\Delta} \quad (29)$$

which can be easily shown to reduce to the Compton wavelength (Cf. also ref.[29]).

Thus the emergence of Benard cell like elementary particles from the Quantum Vacuum mimics the Landau-Ginsburg phase transition. In this case we have a non local growth of correlations reminiscent of the standard inflation theory.

As is known, the interesting aspects of the critical point theory (Cf.ref.[9])

are universality and scale. Broadly, this means that diverse physical phenomena follow the same route at the critical point, on the one hand, and on the other this can happen at different scales, as exemplified for example, by the coarse graining techniques of the Renormalization Group [44]. To highlight this point we note that in critical point phenomena we have the reduced order parameter  $\bar{Q}$  (which gives the fraction of the excess of new states) and the reduced correlation length  $\bar{\xi}$  (which follows from (29)). Near the critical point we have relations [30] like

$$(\bar{Q}) = |t|^\beta, (\bar{\xi}) = |t|^{-\nu}$$

Whence

$$\bar{Q}^\nu = \bar{\xi}^\beta \quad (30)$$

In (30) typically  $\nu \approx 2\beta$ . As  $\bar{Q} \sim \frac{1}{\sqrt{N}}$  because  $\sqrt{N}$  particles are created fluctuationally, given  $N$  particles, and in view of the fractal two dimensionality of the path

$$\bar{Q} \sim \frac{1}{\sqrt{N}}, \bar{\xi} = (l/R)^2 \quad (31)$$

This gives the Eddington formula,

$$R = \sqrt{N}l$$

which is nothing but (6).

There is another way of looking at this. The noncommutative geometry (27) brings out the primacy of the Quantum of Area. Indeed this has been noted from the different perspective of Black Hole Thermodynamics too [45]. We would also like to point out that a similar treatment leads from the Planck scale to the Compton scale.

In other words the creation of particles is the result of a critical point phase transition and subsequent coarse graining (Cf. also ref.[45]).

The above model apart from mimicking inflation also explains as we saw, the so called miraculous large number coincidences.

The peculiarity of these relations as we saw is that they tie up large scale parameters like the radius or age of the universe or the Hubble constant with microphysical parameters like the mass, charge and the Compton scales of an elementary particle and the gravitational constant. That is, the universe appears to have a Machian or holistic feature. One way to understand why the large and the small are tied up is to remember, as we saw in the earlier

chapter, that there is an underpinning of normal mode Planck oscillators, that is, collective phenomena all across the universe.

### 3 The Nature of Space Time

As we noted earlier all of Classical Physics and Quantum Theory, is based on the Minkowski spacetime, as for example in the case of Quantum Field Theory, or Riemannian spacetime as in the case of General Relativity. In the non relativistic theories, Newtonian spacetime, is used, which is a special case of Minkowskian spacetime. But in all these cases the common denominator is that we are dealing with a differentiable manifold.

This breaks down however in Quantum Gravity including the author's approach, String Theory and other approaches, be it at the Planck scale, or at the Compton scale [24, 2, 46, 47]. The underlying reason for this breakdown of a differentiable spacetime manifold is the Uncertainty Principle—as we go down to arbitrarily small spacetime intervals, we encounter arbitrarily large energy momenta. As Wheeler put it [8], “no prediction of spacetime, therefore no meaning for spacetime is the verdict of the Quantum Principle. That object which is central to all of Classical General Relativity, the four dimensional spacetime geometry, simply does not exist, except in a classical approximation.” Before proceeding to analyse the nature of spacetime beyond the classical approximation, let us first analyse briefly the nature of classical spacetime itself.

We can get an insight into the nature of the usual spacetime by considering the well known formulation of Quantum Theory in terms of stochastic processes more precisely, a double Wiener process which, as we saw, models fuzzy spacetime [17, 21].

In the stochastic approach, we deal with a double Wiener process which leads to a complex velocity  $V - iU$ . It is this complex velocity that leads to Quantum Theory from the usual diffusion theory as seen in the previous Chapter. To see this in a simple way, let us write the usual diffusion equation as

$$\Delta x \cdot \Delta x = \frac{h}{m} \Delta t \equiv \nu \Delta t \quad (32)$$

We saw that equation (32) can be rewritten as the usual Quantum Mechanical relation,

$$m \frac{\Delta x}{\Delta t} \cdot \Delta x = h = \Delta p \cdot \Delta x \quad (33)$$

We are dealing here, with phenomena within the Compton or de Broglie wavelength.

We now treat the diffusion constant  $\nu$  to be very small, but non vanishing. That is, we consider the semi classical case. This is because, a purely classical description, does not provide any insight.

It is well known that in this situation we can use the WKB approximation [48]. Whence the right hand side of the wave function,

$$\psi = \sqrt{\rho} e^{i/\hbar S}$$

goes over to, in the one dimensional case, for simplicity,

$$(p_x)^{-\frac{1}{2}} e^{\frac{1}{\hbar} \int p(x) dx}$$

so that we have, on comparison,

$$\rho = \frac{1}{p_x} \quad (34)$$

$\rho$  being the probability density. In this case the condition  $U \approx 0$ , that is, the velocity potential becoming real, implies

$$\nu \cdot \nabla \ln(\sqrt{\rho}) \approx 0 \quad (35)$$

This semi classical analysis suggests that  $\sqrt{\rho}$  is a slowly varying function of  $x$ , infact each of the factors on the left side of (35) would be  $\sim 0(\hbar)$ , so that the left side is  $\sim 0(\hbar^2)$  (which is being neglected). Then from (34) we conclude that  $p_x$  is independent of  $x$ , or is a slowly varying function of  $x$ . The equation of continuity now gives

$$\frac{\partial \rho}{\partial t} + \vec{\nabla}(\rho \vec{v}) = \frac{\partial \rho}{\partial t} = 0$$

That is the probability density  $\rho$  is independent or nearly so, not only of  $x$ , but also of  $t$ . We are thus in a stationary and homogenous scenario. This is strictly speaking, possible only in a single particle universe, or for a completely isolated particle, without any effect of the environment. Under these



circumstances we have the various conservation laws and the time reversible theory, all this taken over into Quantum Mechanics as well. This is an approximation valid for small, incremental changes, as indeed is implicit in the concept of a differentiable space time manifold.

Infact the well known displacement operators of Quantum Theory which define the energy momentum operators are legitimate and further the energy and momenta are on the same footing only under this approximation[49].

We would now like to point out the well known close similarity between the formulation mentioned above and the hydrodynamical formulation for Quantum Mechanics, which also leads to identical equations on writing the wave function as above. These two approaches were reconciled by considering quantized vortices at the Compton scale (Cf.[15]).

To proceed further, we start with the Schrodinger equation

$$i\hbar \frac{\partial \psi}{\partial t} = -\frac{\hbar^2}{2m} \nabla^2 \psi + V\psi \quad (36)$$

Remembering that for momentum eigen states we have, for simplicity, for one dimension

$$\frac{\hbar}{i} \frac{\partial}{\partial x} \psi = p\psi \quad (37)$$

where  $p$  is the momentum or  $p/m$  is the velocity  $v$ , we take the derivative with respect to  $x$  (or  $\vec{x}$ ) of both sides of (36) to obtain, on using (37),

$$i\hbar \frac{\partial (v\psi)}{\partial t} = -\frac{\hbar^2}{2m} \nabla^2 (v\psi) + \frac{\partial V}{\partial x} \psi + Vv\psi \quad (38)$$

We would like to compare (38) with the well known equation for the velocity in hydrodynamics[50, 51], following from the Navier-Stokes equation,

$$\rho \frac{\partial v}{\partial t} = -\nabla p - \rho \alpha T g + \mu \nabla^2 v \quad (39)$$

In (39)  $v$  is a small perturbational velocity in otherwise stationary flow between parallel plates separated by a distance  $d$ ,  $p$  is a small pressure,  $\rho$  is the density of the fluid,  $T$  is the temperature proportional to  $Q(z)v$ ,  $\mu$  is the Navier-Stokes coefficient and  $\alpha$  is the coefficient of volume expansion with temperature. Also required would be

$$\beta \equiv \frac{\Delta T}{d}.$$

$v$  itself is given by

$$v_z = W(z) \exp(\sigma t + \imath k_x x + \imath k_y y), \quad (40)$$

$z$  being the coordinate perpendicular to the fluid flow.

We can now see the parallel between equations (38) and (39). To verify the identification we would require that the dimensionless Rayleigh number

$$R = \frac{\alpha \beta g d^4}{\kappa \nu}$$

should have an analogue in (38) which is dimensionless,  $\kappa, \nu$  being the thermometric conductivity and viscosity.

Remembering that

$$\nu \sim \frac{h}{m}$$

and

$$d \sim \lambda$$

where  $\lambda$  is the Compton wavelength in the above theory (Cf.[52] for details) and further we have

$$\rho \propto f(z)g = V \quad (41)$$

for the identification between the hydrostatic energy and the energy  $V$  of Quantum Mechanics, it is easy using (41) and earlier relations to show that the analogue of  $R$  is

$$(c^2/\lambda^2) \cdot \lambda^4 \cdot (m/h)^2 \quad (42)$$

The expression (42) indeed is dimensionless and of order 1. Thus the mathematical identification is complete.

Before proceeding, let us look at the physical significance of the above considerations (Cf.[53] for a graphic description.) Under conditions of stationery flow, when the difference in the temperature between the two plates is negligible there is total translational symmetry, as in the case of the displacement operators of Quantum Theory. But when there is a small perturbation in the velocity (or equivalently the temperature difference), then beyond a critical value the stationarity and homogeneity of the fluid is disrupted, or the symmetry is broken and we have the phenomena of the formation of Benard cells, which are convective vortices and can be counted. This infact is the "birth" of space It must be stressed that before the formation of the Benard cells, there is no "space", that is, no point to distinguish from or relate to

another point. Only with the formation of the cells are we able to label space points.

In the context of the above identification, the Benard cells would correspond to the formation of “quantized vortices” at the Compton (Planck) scale from the ZPF, as we saw, which latter had been discussed in detail in the literature (Cf.[54]) from the ZPF. This phase transition would correspond to the “formation” of spacetime. As discussed in detail these “quantized vortices” can be identified with elementary particles. Interestingly, as noted Einstein himself considered electrons as condensates from a background electromagnetic field. All this ties up with the discussion in the previous section.

However in order to demonstrate that the above formulation is not a mere mathematical analogy, we have to show that the critical value of the wave number  $k$  in the expression for the velocity in the hydrodynamical flow (40) is the same as the critical length, the Compton length. In terms of the dimensionless wave number  $k' = k/d$ , this critical value is given by[50]

$$k'_c \sim 1$$

In the case of the “quantized vortices” at the Compton scale  $l$ , remembering that  $d$  is identified with  $l$  itself we have,

$$l'_c(\equiv)k'_c \sim 1,$$

exactly as required.

In this connection it may be mentioned that due to fluctuations in the Zero Point Field or the Quantum Vacuum, there would be fluctuations in the metric given by as is known,

$$\Delta g \sim l_P/l$$

where  $l_P$  is the Planck length and  $l$  is a small interval under consideration. At the same time the fluctuation in the curvature of space would be given by

$$\Delta R \sim l_P/l^3$$

Normally these fluctuations are extremely small but as discussed in detail elsewhere[?], this would imply that at the Compton scale of a typical elementary particle  $l \sim 10^{-11}cms$ , the fluctuation in the curvature would be  $\sim 1$ . This is symptomatic of the formation of what we have termed above as elementary particle “quantized vortices”.

Further if a typical time interval between the formation of such “quantized vortices” which are the analogues of the Benard cells is  $\tau$ , in this case the Compton time, then as in the theory of the Brownian Random Walk[78], the mean time extent would be given by

$$T \sim \sqrt{N}\tau \quad (43)$$

where  $N$  is the number of such quantized vortices or elementary particles (Cf.also [15]). This is equation (6) - that is, the equation (43) holds good for the universe itself because  $T$  the age of the universe  $\sim 10^{17}secs$  and  $N$  the number of elementary particles  $\sim 10^{80}$ ,  $\tau$  being the Compton time  $\sim 10^{-23}secs$ . Interestingly, this “phase transition” nature of time would automatically make it irreversible, unlike the conventional model in which time is reversible. We will return to these considerations later in this section. It may be mentioned that an equation similar to (43) can be deduced by the same arguments for space extension also as indeed we did, and this time we get back the well known Eddington formula viz.,

$$R \sim \sqrt{N}l \quad (44)$$

where  $R$  is the extent or radius of the universe and  $l$  is the cell size or Compton wavelength. We can similarly characterize the formation of elementary particles themselves from cells at the Planck scale.

Once we recognize the minimum space time extensions, then we immediately are lead to the underlying non commutative geometry encountered in the earlier chapter and given by equation (27):

$$[x, y] = 0(l^2), [x, p_x] = i\hbar[1 + 0(l^2)], [t, E] = i\hbar[1 + 0(\tau^2)] \quad (45)$$

As we saw relations like (27) are Lorentz invariant. At this stage we recognise the nature of spacetime as given by (27) in contrast to the stationary and homogeneous spacetime discussed earlier. Witten[26, 55] has called this Fermionic spacetime as contrasted with the usual Bosonic spacetime. Indeed we traced the origins of the Dirac equation of the electron to (27). We also argued that (27) provides the long sought after reconciliation between electromagnetism and gravitation[23, 42].

The usual differentiable spacetime geometry can be obtained from (27) if  $l^2$  is neglected, and this is the approximation that has been implicit.

Thus spacetime is a collection of such cells or elementary particles. As

pointed out earlier, this spacetime emerges from a homogeneous stationary non or pre spacetime when the symmetry is broken, through random processes. The question that comes up then is, what is the metric which we use? This has been touched upon earlier, and we will examine it again.

We first make a few preliminary remarks. When we talk of a metric or the distance between two "points" or "particles", a concept that is implicit is that of topological "nearness" - we require an underpinning of a suitably large number of "open" sets[56]. Let us now abandon the absolute or background space time and consider, for simplicity, a universe (or set) that consists solely of two particles. The question of the distance between these particles (quite apart from the question of the observer) becomes meaningless. Indeed, this is so for a universe consisting of a finite number of particles. For, we could isolate any two of them, and the distance between them would have no meaning. We can intuitively appreciate that we would infact need distances of intermediate or more generally, other points.

In earlier work[57, 58], motivated by physical considerations we had considered a series of nested sets or neighbourhoods which were countable and also whose union was a complete Hausdorff space. The Urysohn Theorem was then invoked and it was shown that the space of the subsets was metrizable. Let us examine this in more detail.

Firstly we observe that in the light of the above remarks, the concepts of open sets, connectedness and the like reenter in which case such an isolation of two points would not be possible.

More formally let us define a neighbourhood of a particle (or point or element)  $A$  of a set of particles as a subset which contains  $A$  and atleast one other distinct element. Now, given two particles (or points) or sets of points  $A$  and  $B$ , let us consider a neighbourhood containing both of them,  $n(A, B)$  say. We require a non empty set containing atleast one of  $A$  and  $B$  and atleast one other particle  $C$ , such that  $n(A, B) \subset n(A, C)$ , and so on. Strictly, this "nested" sequence should not terminate. For, if it does, then we end up with a set  $n(A, P)$  consisting of two isolated "particles" or points, and the "distance"  $d(A, P)$  is meaningless.

We now assume the following property[57, 58]: Given two distinct elements (or even subsets)  $A$  and  $B$ , there is a neighbourhood  $N_{A_1}$  such that  $A$  belongs to  $N_{A_1}$ ,  $B$  does not belong to  $N_{A_1}$  and also given any  $N_{A_1}$ , there exists a neighbourhood  $N_{A_{\frac{1}{2}}}$  such that  $A \subset N_{A_{\frac{1}{2}}} \subset N_{A_1}$ , that is there exists an infinite topological closeness.

From here, as in the derivation of Urysohn's lemma[59], we could define a mapping  $f$  such that  $f(A) = 0$  and  $f(B) = 1$  and which takes on all intermediate values. We could now define a metric,  $d(A, B) = |f(A) - f(B)|$ . We could easily verify that this satisfies the properties of a metric.

With the same motivation we will now deduce a similar result, but with different conditions. In the sequel, by a subset we will mean a proper subset, which is also non null, unless specifically mentioned to be so. We will also consider Borel sets, that is the set itself (and its subsets) has a countable covering with subsets. We then follow a pattern similar to that of a Cantor ternary set [56, 60]. So starting with the set  $N$  we consider a subset  $N_1$  which is one of the members of the covering of  $N$  and iterate this process so that  $N_{12}$  denotes a subset belonging to the covering of  $N_1$  and so on.

We note that each element of  $N$  would be contained in one of the series of subsets of a sub cover. For, if we consider the case where the element  $p$  belongs to some  $N_{12\dots m}$  but not to  $N_{1,2,3\dots m+1}$ , this would be impossible because the latter form a cover of the former. In any case as in the derivation of the Cantor set, we can put the above countable series of sub sets of sub covers in a one to one correspondence with suitable sub intervals of a real interval  $(a, b)$ .

### Case I

If  $N_{1,2,3\dots m} \rightarrow$  an element of the set  $N$  as  $m \rightarrow \infty$ , that is if the set is closed, we would be establishing a one to one relationship with points on the interval  $(a, b)$  and hence could use the metric of this latter interval, as seen earlier.

### Case II

It is interesting to consider the case where in the above iterative countable process, the limit does not tend to an element of the set  $N$ , that is set  $N$  is not closed and has what we may call singular points. We could still truncate the process at  $N_{1,2,3\dots m}$  for some  $m > L$  arbitrary and establish a one to one relationship between such truncated subsets and arbitrarily small intervals in  $a, b$ . We could still speak of a metric or distance between two such arbitrarily small intervals.

This case is of interest because we described elementary particles as, what we have called Quantum Mechanical Kerr-Newman Black Holes or vortices, where we have a length of the order of the Compton wavelength as seen in the previous sections, within which spacetime as we know it breaks down. Such cut offs as seen lead to a non commutative geometry (27) and what may be called fuzzy spaces[61, 2].(We note that the centre of the vortex is a singular point). In any case, the number of particles in the universe is of the

order  $10^{80}$ , which approximates infinity from a physicist's point of view. Interestingly, we usually consider two types of infinite sets - those with cardinal number  $n$  corresponding to countable infinities, and those with cardinal number  $c$  corresponding to a continuum, there being nothing in between [60]. This is the well known but unproven Continuum Hypotheses.

What we have shown with the above process is that it is possible to conceive an intermediate possibility with a cardinal number  $n^p, p > 1$ .

In the above considerations three properties are important: the set must be closed i.e. it must contain all its limit points, perfect i.e. in addition each of its points must be a limit point and disconnected i.e. it contains no nonnull open intervals. Only the first was invoked in Case I.

Finally we notice again the holistic feature. A metric emerges by considering large encompassing sets. It may be remarked that much of Quantum Theory, like much of Classical Theory was couched in the concepts of Newtonian two body mechanics and determinism. The moment we consider even a three body problem, as was realized by Poincare more than a century ago, the picture gets altered. As he noted [62], "If we knew exactly the laws of nature and the situation of the universe at the initial moment, we could predict exactly the situation of that same universe at a succeeding moment. But even if it were the case that the natural laws had no longer any secret for us, we could still know the situation approximately. If that enabled us to predict the succeeding situation with the same approximation, that is all we require, and we should say that the phenomenon had been predicted, that it is governed by the laws. But it is not always so; it may happen that small differences in the initial conditions produce very great ones in the final phenomena. A small error in the former will produce an enormous error in the latter. Prediction becomes impossible." In a similar vein, Prigogine observes [53], "Our physical world is no longer symbolized by the stable and periodic planetary motions that are at the heart of classical mechanics. It is a world of instabilities and fluctuations..."

Indeed, the departure from the two body formulation began with electromagnetism itself, which has to invoke the environment.

We now return to the current view of Planck scale oscillators in the background dark energy or Quantum Vacuum. In this context we saw that elementary particles can be considered to be normal modes of  $n \sim 10^{40}$  Planck oscillators in the ground state, while the entire universe itself has an underpinning of  $\bar{N} \sim 10^{120}$  such oscillators, there being  $N \sim 10^{80}$  elementary particles in the universe [32, 43]. These Planck oscillators are formed out

of the Quantum Vacuum (or dark energy). Thus we have,  $m_P c^2$  being the energy of each Planck oscillator,  $m_P$  being the Planck mass,  $\sim 10^{-5} gms$ ,

$$m = \frac{m_P}{\sqrt{n}} \quad (46)$$

$$l = \sqrt{n} l_P, \tau = \sqrt{n} \tau_P, n = \sqrt{N} \quad (47)$$

where  $m$  is the mass of a typical elementary particle, taken to be a pion in the literature. The ground state of  $\bar{N}$  such Planck oscillators would be, in analogy to (46),

$$\bar{m} = \frac{m_P}{\sqrt{N}} \sim 10^{-65} gms \quad (48)$$

The universe is an excited state and consists of  $N$  such ground state levels and so we have, from (48)

$$M = \bar{m} N = \sqrt{N} m_P \sim 10^{55} gms,$$

as required,  $M$  being the mass of the universe.

Due to the fluctuation  $\sim \sqrt{n}$  in the levels of the  $n$  oscillators making up an elementary particle, the energy is, remembering that  $mc^2$  is the ground state,

$$\Delta E \sim \sqrt{n} mc^2 = m_P c^2,$$

using (47), and so the indeterminacy time is

$$\frac{\hbar}{\Delta E} = \frac{\hbar}{m_P c^2} = \tau_P,$$

as indeed we would expect.

The corresponding minimum indeterminacy length would therefore be  $l_P$ . One of the consequences of the minimum spacetime cut off as we saw is that the Heisenberg Uncertainty Principle takes an extra term as we saw previously [63]. Thus as we saw

$$\Delta x \approx \frac{\hbar}{\Delta p} + \alpha \frac{\Delta p}{\hbar}, \alpha = l^2 (\text{or } l_P^2) \quad (49)$$

where  $l$  (or  $l_P$ ) is the minimum interval under consideration. This is just (28). The first term gives the usual Heisenberg Uncertainty Principle.

Application of the time analogue of (49) for the indeterminacy time  $\Delta t$  for the



fluctuation in energy  $\Delta\bar{E} = \sqrt{N}mc^2$  in the  $N$  particle states of the universe gives exactly as above,

$$\Delta t = \frac{\Delta E}{\hbar} \tau_P^2 = \frac{\sqrt{N}mc^2}{\hbar} \tau_P^2 = \frac{\sqrt{N}m_P c^2}{\sqrt{n}\hbar} \tau_P^2 = \sqrt{n} \tau_P = \tau,$$

wherein we have used (47). In other words, for the fluctuation  $\sqrt{N}$ , the time is  $\tau$ . It must be re-emphasized that the Compton time  $\tau$  of an elementary particle, is an interval within which there are unphysical effects like zitterbewegung - as pointed out by Dirac, it is only on averaging over this interval, that we return to meaningful Physics. This gives us,

$$dN/dt = \sqrt{N}/\tau \quad (50)$$

Equation (78) is identical to (3), the starting point for the cosmology discussed. Here we have derived this relation from a consideration of the underlying Planck oscillators. On the other hand due to the fluctuation in the  $\sqrt{N}$  oscillators constituting the universe, the fluctuational energy is similarly given by

$$\sqrt{N}\bar{m}c^2,$$

which is the same as (48) above. Another way of deriving (50) is to observe that as  $\sqrt{n}$  particles appear fluctuationally in time  $\tau_P$  which is, in the elementary particle time scales,  $\sqrt{n}\sqrt{n} = \sqrt{N}$  particles in  $\sqrt{n}\tau_P = \tau$ . That is, the rate of the fluctuational appearance of particles is

$$\left( \frac{\sqrt{n}}{\tau_P} \right) = \frac{\sqrt{N}}{\tau} = dN/dt$$

which is (50). From here by integration,

$$T = \sqrt{N}\tau$$

$T$  is the time elapsed from  $N = 1$  and  $\tau$  is the Compton time. This gives  $T$  its origin in the fluctuations - there is no smooth “background” (or “being”) time - the root of time is in “becoming”. It is the time of a Brownian double Wiener process: A step  $l$  gives a step in time  $l/c \equiv \tau$  and therefore  $\Delta x = \sqrt{N}l$  gives  $T = \sqrt{N}\tau$ . Time is born out of acausal fluctuations which are random and therefore irreversible. Indeed, there is no background time.

Time is proportional to  $\sqrt{N}$ ,  $N$  being the number of particles which are being created spontaneously from the ZPF.

The time we use is what may be called Stationary time and it is an approximation as we saw [21]. Further, Quantum Mechanics, Gravitation etc. follow from here. In Quantum Mechanics, the measurement of the observer triggers the acausal collapse of the wave function - an irreversible event - but the wave function itself satisfies a deterministic and reversible equation paradoxically. Yet the universe is “irreversible”. It appears spontaneous irreversibility or indeterministic time [15] is the real time. This can be contrasted to the usual time reversible Quantum Theory.

We observe that [65]

$$\psi(r, t) = \frac{1}{(2\pi\hbar)^{3/2}} \int a(p) \exp \left[ \frac{i}{\hbar} \left( p \cdot r - \frac{p^2}{2m} t \right) \right] dp,$$

$a(p)$  being independent of time. So we have at any other time  $t'$ :

$$a(p) = \frac{1}{(2\pi\hbar)^{3/2}} \int \psi(r', t') \exp \left[ -\frac{i}{\hbar} \left( p \cdot r' - \frac{p^2}{2m} t' \right) \right] dr'$$

Substitution yields the result

$$\psi(r, t) = \int K(r, t; r', t') \psi(r', t') dr', \quad (51)$$

the Kernel function  $K$  being given by

$$K(r, t; r', t') = \frac{1}{(2\pi\hbar)^3} \int \exp \left\{ \frac{i}{\hbar} \left[ p \cdot (r - r') - \frac{p^2}{2m} (t - t') \right] \right\} dp$$

or after some manipulation, in the form

$$K(r, t; r', t') = \left[ \frac{2\pi i \hbar}{m} (t - t') \right]^{-3/2} \exp \left[ i \frac{m}{2\hbar} \frac{|r - r'|^2}{(t - t')} \right]$$

The point is that in (51)  $\psi(r, t)$  at  $t$  is given in terms of a linear expansion of  $\psi(r, t')$  at earlier times  $t'$ . But what is to be noted is, the symmetry between  $t$  and  $t'$ . This is not surprising as the original Schrodinger equation remains unchanged under  $t \rightarrow -t$ .

Thus it is possible to understand the fluctuations encountered in Section 2,

that is, the equation (78) which was the starting point for fluctuational energy in terms of the underpinning of Planck scale oscillators in the Quantum Vacuum.

We would now like to make some remarks. Starting from a completely different point of view namely Black Hole Thermodynamics, Landsberg [64] deduced that the smallest observable mass in the universe is  $\sim 10^{-65}gms$ , which is exactly the minimum mass given in (48).

Further due to the fluctuational appearance of  $\sqrt{N}$  particles, the fluctuational mass associated with each of the  $N$  particles in the universe is

$$\frac{\sqrt{N}m}{N} = \frac{m}{\sqrt{N}} \sim 10^{-65}gms,$$

that is once again the smallest observable mass or ground state mass in the universe.

## 4 Further Considerations

1. We will now provide yet another rationale for (3). Let us start with equations encountered earlier,

$$R = \sqrt{N}l$$

$$\frac{Gm^2}{e^2} = \frac{1}{\sqrt{N}} \sim 10^{-40}$$

or the Weinberg formula

$$m = \left( \frac{H\hbar^2}{Gc} \right)^{\frac{1}{3}}$$

where  $N \sim 10^{80}$  is the number of elementary particles, typically pions, in the universe. On the other hand the ratio of the electromagnetic and Gravitational coupling constants, is deducible from (3). The very mysterious feature of the 'Weinberg' formula was stressed by Weinberg himself "...it should be noted that the particular combination of  $\hbar, H, G$ , and  $c$  appearing (in the formula) is very much closer to a typical elementary particle mass than other random combinations of these quantities....

In contrast, (the formula) relates a single cosmological parameter,  $H$ , to the

fundamental constants  $\hbar, G, c$  and  $m$ , and is so far unexplained...”

Relations like (9) and (8) inspired the Dirac Large Number Cosmology. All these relations are to be taken in the order of magnitude sense.

We will now take a different route and provide an alternative theoretical rationale for equations (71), (8) and (9), and in the process light will be shed on the new cosmological model and the nature of gravitation.

Following Sivaram [66] we consider the gravitational self energy of the pion. This is given by

$$\frac{Gm^2}{l} = Gm^2/(\hbar/mc)$$

If this energy were to have a life time of the order of the age of the universe,  $T$ , then we have by the Uncertainty relation

$$\left(\frac{Gm^3c}{\hbar}\right)(T) \approx \hbar \quad (52)$$

As  $T = \frac{1}{H}$ , this immediately gives us the Weinberg formula (7). It must be observed again that (52) gives a time dependent gravitational constant  $G$ . We could also derive (7) by using a relation given by Landsberg [65]. We use the fact that the mass of a particle is given by

$$m(b) \sim \left(\frac{\hbar^3 H}{G^2}\right)^{1/5} \left(\frac{c^5}{\hbar H^2 G}\right)^{b/15} \quad (53)$$

where  $b$  is an unidentified constant. Whence we have

$$m(b) \sim G^{-3/5} G^{-3b/15} = G^{-(b+1)/5}$$

The mass that would be time independent, if  $G$  were time dependent would be given by the value

$$b = -1$$

With this value of  $b$  (53) gives back (7).

Let us now proceed along a different track. We rewrite (52) as

$$G = \frac{\hbar^2}{m^3 c} \cdot \frac{1}{T} \quad (54)$$

If we use the fact that  $R = cT$ , then (54) can be written as

$$G = \frac{\hbar^2}{m^3 R} \quad (55)$$

Let us now use the well known relation encountered earlier [67]

$$R = \frac{GM}{c^2}, \quad (56)$$

There are several derivations of (56). For example in a uniformly expanding Friedman universe, we have

$$\dot{R}^2 = \frac{8\pi G\rho R^2}{3}$$

If we substitute the value  $\dot{R} = c$  at the radius of the universe, then we recover (56). If we use (56) in (55) we will get

$$G^2 = \frac{\hbar^2 c^2}{m^3 M} \quad (57)$$

Let  $M/m = N$  be called the number of elementary particles in the universe. Infact this is just (1). Then (57) can be written as (10),

$$G = \frac{\hbar c}{m^2 \sqrt{N}}$$

which can also be written as (8)

$$Gm^2/e^2 \sim \frac{1}{\sqrt{N}}$$

Whence we get (9)

$$\sqrt{N}l = R$$

We now remark that (54) shows an inverse dependence on time of the gravitation constant, while (10) shows an inverse dependence on  $\sqrt{N}$ . Equating the two, we get back,

$$T = \sqrt{N}\tau$$

the relation (4) which we have encountered several times. If we now take the time derivative of (10) and use (4), we get (3)

$$\dot{N} = \frac{\sqrt{N}}{\tau}$$

This equation is the same as (78) or (3). To put it briefly in a phase transition from the Quantum Vacuum  $\sqrt{N}$  particles appear within the Compton time

$\tau$ . In terms of our unidirectional concept of time, we could say that particles appear and disappear, but the nett result is the appearance of  $\sqrt{N}$  particles. We now make a few remarks. Firstly it is interesting to note that  $\sqrt{N}m$  will be the mass added to the universe. Let us now apply the well known Beckenstein formula for the life time of a mass  $M$  viz., [67],

$$t \approx G^2 M^3 / \hbar c^4$$

to the above mass. The life time as can be easily verified turns out to be exactly the age of the universe!

A final remark. To appreciate the role of fluctuations in the otherwise mysterious Large Number relations, let us follow Hayakawa and consider the excess of electric energy due to the fluctuation  $\sim \sqrt{N}$  of the elementary particles in the universe and equate it to the inertial energy of an elementary particle. We get

$$\frac{\sqrt{N}e^2}{R} = mc^2$$

This gives us back (8) if we use (56). If we use (9) on the other hand, we get

$$e^2/mc^2 = l,$$

another well known relation from micro physics.

2. We note that as is well known, a background ZPF of the kind we have been considering can explain the Quantum Mechanical spin half as also the anomalous  $g = 2$  factor for an otherwise purely classical electron [68, 69, 70]. The key point here is (Cf.ref.[68]) that the classical angular momentum  $\vec{r} \times m\vec{v}$  does not satisfy the Quantum Mechanical commutation rule for the angular momentum  $\vec{J}$ . However when we introduce the background Zero Point Field, the momentum now becomes

$$\vec{J} = \vec{r} \times m = \vec{v} + (e/2c)\vec{r} \times (\vec{B} \times \vec{r}) + (e/c)\vec{r} \times \vec{A}^0, \quad (58)$$

where  $\vec{A}^0$  is the vector potential associated with the ZPF and  $\vec{B}$  is an external magnetic field introduced merely for convenience, and which can be made vanishingly small.

It can be shown that  $\vec{J}$  in (58) satisfies the Quantum Mechanical commutation relation for  $\vec{J} \times \vec{J}$ . At the same time we can deduce from (58)

$$\langle J_z \rangle = -\frac{1}{2}\hbar\omega_0/|\omega_0| \quad (59)$$

Relation (59) gives the correct Quantum Mechanical results referred to above. From (58) we can also deduce that

$$l = \langle r^2 \rangle^{\frac{1}{2}} = \left( \frac{\hbar}{mc} \right) \quad (60)$$

After several fruitless decades of attempts to provide a unified description of Gravitation and Electromagnetism, it has been realized that differentiable space time has to be abandoned in favour of one with a minimum scale. It is now generally accepted that the Planck scale defines the minimum scale for the universe [3, 2, 4, 71, 43, 32]. In these schemes there is a maximal mass, the Planck mass  $m_P \sim 10^{-5} gms$  which is defined by

$$m_P = \left( \frac{\hbar c}{G} \right)^{1/2} \approx 10^{-5} gms \quad (61)$$

Using the value for  $m_P$  we can define the Planck length  $l_P \sim 10^{-33} cms$  and the Planck time  $t_P \sim 10^{-42} secs$ , which are the Compton lengths and times for the mass in (61). It may be mentioned that these values were postulated by Planck himself. Today the values for the minimum scale as given in (61) are taken for granted. We first provide a rationale for the numerical value of the Planck scale.

## 5 The Planck Scale

Our starting point is the model for the underpinning at the Planck scale for the universe. This is a collection of  $N$  Planck scale oscillators (Cf.refs.[43, 32, 72, 73, 74] for details). We do not need to specify  $N$ . We have in this case the following well known relations

$$R = \sqrt{N}l, K l^2 = kT, \quad \omega_{max}^2 = \frac{K}{m} = \frac{kT}{m l^2} \quad (62)$$

In (62),  $R$  is of the order of the diameter of the universe,  $K$  is the analogue of the spring constant,  $T$  is the effective temperature while  $l$  is the analogue of the Planck length,  $m$  the analogue of the Planck mass and  $\omega_{max}$  is the

frequency-the reason for the subscript *max* will be seen below. We do not yet give  $l$  and  $m$  their usual values as given in (61) for example, but rather try to deduce these values.

We now use the well known result that the individual minimal oscillator particles are black holes or mini universes as shown by Rosen [7]. So using the well known Beckenstein temperature formula for these primordial black holes [67], that is

$$kT = \frac{\hbar c^3}{8\pi Gm}$$

in (62) we get,

$$Gm^2 \sim \hbar c \quad (63)$$

which is another form of (61). We can easily verify that (63) leads to the value  $m \sim 10^{-5} gms$ . In deducing (63) we have used the typical expressions for the frequency as the inverse of the time - the Compton time in this case and similarly the expression for the Compton length. However it must be reiterated that no specific values for  $l$  or  $m$  were considered in the deduction of (63).

We now make two interesting comments. Cercignani and co-workers have shown [75, 76] that when the gravitational energy becomes of the order of the electromagnetic energy in the case of the Zero Point oscillators, that is

$$\frac{G\hbar^2\omega^3}{c^5} \sim \hbar\omega \quad (64)$$

then this defines a threshold frequency  $\omega_{max}$  above in which the oscillations become chaotic.

Secondly from the parallel but unrelated theory of phonons [77, 78], which are also bosonic oscillators, we deduce a maximal frequency given by

$$\omega_{max}^2 = \frac{c^2}{l^2} \quad (65)$$

In (65)  $c$  is, in the particular case of phonons, the velocity of propagation, that is the velocity of sound, whereas in our case this velocity is that of light. Frequencies greater than  $\omega_{max}$  in (65) are meaningless. We can easily verify that (64) and (65) give back (63).

Finally we can see from (62) that, given the value of  $l_P$  and using the value of the radius of the universe, viz.,  $R \sim 10^{27}$ , we can deduce that,

$$N' \sim 10^{120} \quad (66)$$



In a sense the relation (63) can be interpreted in a slightly different vein as representing the scale at which all energy- gravitational and electromagnetic becomes one.

## 6 The Gauge Hierarchy Problem

A long standing puzzle has been the so called Gauge Hierarchy Problem. This deals with the fact that the Planck mass is some  $10^{20}$  times the mass of an elementary particle, for example Gauge Bosons or Protons or Electrons (in the large number sense). Why is there such a huge gap?

We now recall that as already shown we have [10, 11, 79, 72]

$$G = \frac{\hbar c}{m^2 \sqrt{N}} \quad (67)$$

In (67)  $N \sim 10^{80}$  is the well known number of elementary particles in the universe, which features in the Weyl-Eddington relations as also the Dirac Cosmology.

What is interesting about (67) is that it shows gravitation as a distributional effect over all the  $N$  particles in the universe [79, 72].

Let us rewrite (63) in the form

$$G \approx \frac{\hbar c}{m_P^2} \quad (68)$$

remembering that the Planck length is also the Compton length of the Planck mass. (Interestingly an equation like (??) or (4) also follows from Sakharov's treatment of gravitation [80].) A division of (3) and (4) yields

$$m_P^2 = \sqrt{N} m^2 \quad (69)$$

Equation (69) immediately gives the ratio  $\sim 10^{20}$  between the Planck mass and the mass of an elementary particle.

Comparing (67) and (68) we can see that the latter is the analogue of the former in the case  $N \sim 1$ . So while the Planck mass in the spirit of Rosen's isolated universe and the Schwarzschild black hole uses the gravitational interaction in isolation, as seen from (67), elementary particles are involved in the gravitational interaction with all the remaining particles in the universe.

Finally rememebring that  $Gm_P^2 \sim e^2$ , as can also be seen from (68), we get from (67)

$$\frac{e^2}{Gm^2} \sim \frac{1}{\sqrt{N}} \quad (70)$$

Equation (70) is the otherwise empirically well known electromagnetism-gravitation coupling constant ratio, but here it is deduced from the theory. It may be remarked that one could attempt an explanation of (69) from the point of view of SuperSymmetry or Brane theory, but these latter have as yet no experimental validation [81]. On the other hand, the crucial equation (67) was actually part of a predicted dark energy driven accelerating universe with a small cosmological constant (besides a deduction of hitherto empirical Large Number relations) (Cf. for example ref.[10])– all this got dramatic observational confirmation through the works of Perlmutter, Kirshner and others, shortly thereafter.

## References

- [1] Sidharth, B.G., Annales Fondation L. De Broglie, 29 (3), 2004, 1.
- [2] Kempf, A., in “From the Planck length to the Hubble radius”, Ed. A. Zichichi, World Scientific, Singapore, 2000, pp.613ff.
- [3] Veneziano, G., in “The Geometric Universe”, Ed. by S.A. Huggett, et al., Oxford University Press, Oxford, 1998, p.235ff.
- [4] Garay, L.J., Int.J.Mod.Phys.A., 1995, 10 (2), 145-165.
- [5] Sivaram, C., Am.J.Phys., 51(3), 1983, p.277.
- [6] Sidharth, B.G., “The Chaotic Universe: From the Planck to the Hubble Scale”, Nova Science Publishers, Inc., New York, 2001.
- [7] Rosen, N., Int.J.Th.Phys., 32(8), 1993, p.1435-1440.
- [8] Sidharth, B.G., and Popova, A.D., Differential Equations and Dynamical Systems, 4 (3/4), 1996, 431-440.
- [9] Narlikar, J.V., Int.J.Th.Phys., 1983, pp.311-323.
- [10] Sidharth, B.G., Int.J.Mod.Phys.A, 13 (15), 1998, p.2599ff.

- [11] Sidharth, B.G., *Int.J.Th.Phys.*, 37 (4), 1998, p.1307ff.
- [12] Sidharth, B.G., *Il Nuovo Cimento*, 115B (12), (2), 2000, pg.151.
- [13] Norman, E.G., *Am.J.Phys.*, 544, 317, 1986.
- [14] Uzan, J.P., *Rev.Mod.Phys.*, 75, (April 2003), pp.403-455.
- [15] Sidharth, B.G., *Chaos, Solitons and Fractals*, 12(1), 2001, 173-178.
- [16] Sidharth, B.G., *Chaos, Solitons and Fractals*, 13, 2002, pp.1325-1330.
- [17] Nelson, E., *Physical Review*, Vol.150, No.4, October 1966, p.1079-1085.
- [18] Nottale, L., “Fractal Space-Time and Microphysics: Towards a Theory of Scale Relativity”, World Scientific, Singapore, 1993, p.312.
- [19] Smolin, L., in “Quantum Concepts in Space and Time”, Eds., R. Penrose and C.J. Isham, OUP, Oxford, 1986, pp.147-181.
- [20] Kyprianidis, A., *Int.J.Th.Phys.*, 1992, pp.1449-1483.
- [21] Sidharth, B.G., “Concise Encyclopaedia of SuperSymmetry and Non Commutative Structures in Mathematics and Physics”, Eds. J. Bagger, S. Duply, W. Sugel, New York, Kluwer Academic Publishers, 2001.
- [22] Ne’eman, Y., in *Proceedings of the First International Symposium, “Frontiers of Fundamental Physics”*, Eds. B.G. Sidharth and A. Burinskii, Universities Press, Hyderabad, 1999, pp.83ff.
- [23] Sidharth, B.G., *Nuovo Cimento*, 116B (6), 2001, p.4ff.
- [24] Amati, D., in “Sakharov Memorial Lectures”, Eds. L.V. Kaddysch and N.Y. Feinberg, Nova Science, New York, 1992, pp.455ff.
- [25] Guida, R., et al., *Mod.Phys.Lett. A*, Vol.6, No.16, 1991, p.1487-1503.
- [26] Witten, W., *Physics Today*, April 1996, pp.24-30.
- [27] Maggiore, M., *Phys.Lett.B.*, 319, 1993, p.83-86.
- [28] Maggiore, M., *Phys.Rev.D.*, Vol.49, No.10, 1994, p.5182-5187.
- [29] Doplicher, S., et al., *Phys.Lett., B*, 331, 1994, p.39-44.

- [30] Sivaram, C., Am.J.Phys., 50(3), 1982, p.279ff.
- [31] Van Dam, H., et al., Mod.Phys.Lett.,A, Vol.9, No.4, 1994, 335-340.
- [32] Sidharth, B.G., Found.Phys.Lett., 15 (6), 2002, pp.577-583.
- [33] Goodstein, D.L., "States of Matter", Dover Publications Inc., New York, 1985, p.160ff.
- [34] Sidharth, B.G., Proceedings of the Fifth International Symposium on "Frontiers of Fundamental Physics", Universities Press, Hyderabad, 2004 (In Press).
- [35] Sidharth, B.G., Chaos, Solitons and Fractals, 13, 2002, p.1369-1370.
- [36] Sidharth, B.G., Frontiers of Fundamental Physics 4, Plenum Publishers/Kluwer Academic, New York, 2001, p.97-107.
- [37] Kalyana Rama, S., Phys.Lett. B, 519, 2001, p.103-110.
- [38] Moffat, J.W., Int.J.Mod.Phys.,D 2, 1993, p.351, J.W. Moffat, Found.Phys. 23, 1993.
- [39] Magueijo, J., Phys.Rev.D., 62, 2000, 103521.
- [40] Hoyle, F., and Narlikar, J.V., The Astrophysical Journal, 410, June 20 1993, p.437-457.
- [41] Sidharth, B.G., in Instantaneous Action at a Distance in Modern Physics: "Pro and Contra", Eds., A.E. Chubykalo et. al., Nova Science Publishing, New York, 1999.
- [42] Sidharth, B.G., Annales de la Fondation Louis de Broglie, 27 (2), 2002, pp.333ff.
- [43] Sidharth, B.G., Found.Phys.Lett., 17 (5), 2004, pp.503-506.
- [44] Wilson, K.G., Rev.Mod.Phys., 1983, 55, pp.583ff.
- [45] Baez, J., Nature, Vol.421, February 2003, pp.702-703.
- [46] Sidharth, B.G., Found.Phys.Lett., August 2002.

- [47] Madore, J., *Class.Quantum Grav.* 9, 1992, p.69-87.
- [48] Schiff, L.I., "Quantum Mechanics", McGraw Hill, London, 1968.
- [49] Davydov, A.S., "Quantum Mechanics, Pergamon Press, Oxford, 1965, p.655.
- [50] Rai Choudhuri, A., "The Physics of Fluids and Plasma", Cambridge University Press, 1999, pp.137.
- [51] Batchelor, G.K., "Fluid Mechanics", Cambridge University Press, New Delhi, 1993.
- [52] Sidharth, B.G., *Ind.J. of Pure and Applied Physics*, 35, 1997, p.456ff.
- [53] Nicolis, G., and Prigogine, I., "Exploring Complexity", W.H. Freeman, New York, 1989, p.10.
- [54] Vasudevan, R., "Hydrodynamical Formulation of Quantum Mechanics", in "Perspectives in Theoretical Nuclear Physics", Ed.Srinivas Rao, K., and Satpathy, L., Wiley Eastern, New Delhi, 1994, pp.216ff.
- [55] Sidharth, B.G., *Chaos, Solitons and Fractals*, 13 (2), 2002, pp.189-193.
- [56] Simmons, G.F., "Introduction to Topology and Modern Analysis", McGraw Hill Book Co. Inc., New York, 1965, p.135.
- [57] Altaisky, M.V., and Sidharth, B.G., *Chaos, Solitons & Fractals*, Vol. 10, No.2-3, 1999, pp.167-176.
- [58] Sidharth, B.G., *Chaos, Solitons and Fractals*, 14, 2002, p.525-527.
- [59] Munkres, J.R., "Topology", Prentice Hall India, New Delhi, 1988.
- [60] Gullick, D., "Encounters With Chaos", McGraw Hill, New York, 1997, p.114ff.
- [61] Sidharth, B.G., *Gravitation and Cosmology*, 4 (2) (14), 1998, p.158ff.
- [62] Gleick, J., "Chaos: Making a New Science", Viking, New York, 1987, p.321.

- [63] Sidharth, B.G., Chaos, Solitons and Fractals, 15, 2003, 593-595, and several references therein.
- [64] Powell, J.L., and Craseman, B., "Quantum Mechanics", Addison-Wesley, Cambridge, 1961.
- [65] Landsberg, P.T., Am.J.Phys., 51, 1983, pp.274-275.
- [66] Sivaram, C., Astrophysics and Space Science, 88, 1982, p.507-510.
- [67] Zang, L.Z., and Ruffini, R., "Basic Concepts in Relativistic Astrophysics", World Scientific, Singapore, 1983, p.123.
- [68] Sachidanandam, S., Physics Letters, Vol.97A, No.8, 19 September 1983, p.323-324.
- [69] Boyer, T.H., Phys.Rev., 174, 1631, 1968.
- [70] Milonni, P.W., "The Quantum Vacuum: An Introduction to Quantum Electrodynamics", Academic Press, San Diego, 1994.
- [71] Crane, L., and Smolin, L., Nucl.Phys., B267, 1986, p.714.
- [72] Sidharth, B.G., "The Universe of Fluctuations", Springer, Berlin, 2005.
- [73] Sidharth, B.G., (Ed.) "The Great Ideas of Physics", Springer, in press.
- [74] Jack Ng, Y., and Van Dam, H., Mod.Phys.Lett.A., 9 (4), 1994, p.335-340.
- [75] Cercignani, C., Found.Phys.Lett., Vol.11, No.2, 1998, 189-199.
- [76] Cercignani, C., Galgani, L., and Scotti, A., Phys.Lett. 38A, 1972, 403.
- [77] Huang, K., "Statistical Mechanics", Wiley Eastern, New Delhi 1975, pp.75ff.
- [78] Reif, F., "Fundamentals of Statistical and Thermal Physics", McGraw-Hill Book Co., Singapore, 1965.
- [79] B.G. Sidharth, "A Note on the Characterization of Gravitation", Foundation of Physics Letters, May, 2005.

- [80] A.D. Sakharov, Soviet Physics - Doklady, Vol.12, No.11, 1968, p.1040-1041.
- [81] E. Golfain, Chaos, Solitons and Fractals, 22 (3), 2004, pp.513-520.